

Donaldson invariants for non-simply connected manifolds

Marcos Mariño and Gregory Moore

Department of Physics, Yale University

New Haven, CT 06520

marino@genesis5.physics.yale.edu

moore@castalia.physics.yale.edu

We study Coulomb branch (“u-plane”) integrals for $\mathcal{N} = 2$ supersymmetric $SU(2)$, $SO(3)$ Yang-Mills theory on 4-manifolds X of $b_1(X) > 0$, $b_2^+(X) = 1$. Using wall-crossing arguments we derive expressions for the Donaldson invariants for manifolds with $b_1(X) > 0$, $b_2^+(X) > 0$. Explicit expressions for $X = \mathbb{C}P^1 \times F_g$, where F_g is a Riemann surface of genus g are obtained using Kronecker’s double series identity. The result might be useful in future studies of quantum cohomology.

April 11, 1998

1. Introduction

The Donaldson invariants of four-manifolds have been a source of fascination both in mathematics and in physics. While there has been much progress in understanding these invariants, there is more to learn, particularly in terms of the relation of the invariants to Floer homology [1] and to Gromov-Witten invariants [2][3]. Understanding the invariants in these contexts leads to the need to understand the Donaldson invariants for non-simply connected four-manifolds X . Most investigations of Donaldson invariants have focussed on the case $\pi_1(X) = 0$. There exists a mathematical definition in the non-simply connected case [4] but comparatively little is known about this case. This paper derives some new results on Donaldson invariants for 4-manifolds with first Betti number $b_1(X) > 0$. We do not consider the effects of torsion in $H_*(X; \mathbb{Z})$, nor the effects of a nonabelian fundamental group.

Recently, using Witten's physical approach to Donaldson theory [5][6][7][8], a fairly systematic (physical) procedure has been developed for deriving various properties of the Donaldson invariants, including wall-crossing and blowup formulae, and the relation to Seiberg-Witten invariants [9][10][11][12][13]. The systematic procedure, which begins with certain integrals over the Coulomb branch of vacua of an $\mathcal{N} = 2$ SYM theory can be extended to higher rank gauge groups and to nonsimply connected manifolds. In [9][10] partial results were obtained for nonsimply connected manifolds. In this paper a more complete treatment is given for the rank one groups $SU(2), SO(3)$.

Our main results are:

1. A wall-crossing formula for the Donaldson invariants in equations (3.4) and (3.16) below.
2. An expression for the Donaldson invariants in terms of SW invariants. For manifolds of simple type this is given in equation (4.17) below. It is a natural generalization of Witten's formula [7], obtained in the simply connected case.
3. Explicit expressions for the Donaldson invariants for $X = \mathbb{CP}^1 \times F_g$ where F_g is a Riemann surface of genus g . We give answers valid in *both* the chambers $\text{vol}(\mathbb{CP}^1) \rightarrow 0$ and $\text{vol}(F_g) \rightarrow 0$.

The expressions in equations (5.18), (5.20) below might prove useful in future studies of the Gromov-Witten invariants of the moduli space of flat connections on F_g .

2. The u -plane integral for $b_1 > 0$.

In this section we extend and elaborate on the results of section 10 of [9]. We consider an arbitrary insertion of observables, using the proposal for the contact terms in [10].

Consider an $\mathcal{N} = 2$ $SU(2)$ or $SO(3)$ supersymmetric Yang-Mills theory¹ on a compact oriented 4-manifold X of $b_1(X) > 0$. As explained in [9] the Donaldson-Witten generating function can be written as $Z_{DW} = Z_u + Z_{SW}$ where Z_u is the Coulomb branch integral and Z_{SW} is the contribution of the Seiberg-Witten invariants. The Coulomb branch integral is only nonvanishing for $b_2^+(X) = 1$, but, by a procedure explained in [9][12] can be taken as the starting point for a systematic derivation of Z_{DW} . In the case of $b_1(X) > 0$ the Coulomb integral can be obtained by a simple generalization of the arguments in section three of [9]. First of all, the photon partition function includes [8] an integration over b_1 zero modes of the gauge field corresponding to flat connections. These zero modes span the tangent space to a torus of dimension b_1 , $\mathbf{T}^{b_1} = H^1(X, \mathbb{R})/H^1(X, \mathbb{Z})$. The zero modes of the one-forms ψ live in this tangent space. As a consequence of having these extra zero modes, the photon partition function is

$$(\text{Im}\tau)^{\frac{1}{2}(b_1-1)}\theta_0(\tau, \bar{\tau}), \quad (2.1)$$

where $\theta_0(\tau, \bar{\tau})$ is the Siegel-Narain theta function introduced in [8][14][9].

The next ingredient comes from the measure for the ψ -fields. The expansion in zero modes reads $\psi = \sum_{i=1}^{b_1} c_i \beta_i$, where β_i , $i = 1, \dots, b_1$ is an integral basis of harmonic one-forms, and we identify $\mathcal{H}^1(X) \simeq H^1(X, \mathbb{Z})$. The c_i are Grassmann variables. The measure for the ψ fields is then:

$$\prod_{i=1}^{b_1} \frac{dc_i}{(\text{Im}\tau)^{\frac{1}{2}}} = (\text{Im}\tau)^{-\frac{b_1}{2}} \prod_{i=1}^{b_1} dc_i. \quad (2.2)$$

We can consider the c_i as a basis of one -forms $\beta_i^\sharp \in H^1(\mathbf{T}^{b_1}, \mathbb{Z})$, dual to $\beta_i \in H^1(X, \mathbb{Z})$. In this way we can identify

$$\psi = \sum_{i=1}^{b_1} \beta_i \otimes \beta_i^\sharp = c_1(\mathcal{L}), \quad (2.3)$$

where $\mathcal{L}$ is the universal flat line bundle over $\mathbf{T}^{b_1} \times X$.

¹ For simplicity we do not include hypermultiplets.

Taking into account (2.1) and (2.2), we see that the Coulomb integral (without any insertion of observables) can be written as [9]:

$$\begin{aligned}
Z_u = & 2 \int [da \, d\bar{a} \, d\eta \, d\chi] \int_{Pic(X)} d\psi \int dDA^\times B^\sigma y^{-1/2} \\
& \exp \left[\frac{1}{8\pi} \int (\text{Im}\tau) D \wedge *D \right] \exp \left[-i\pi\bar{\tau}\lambda_+^2 - i\pi\tau\lambda_-^2 + \pi i(\lambda, w_2(X)) \right] \\
& \exp \left[-\frac{i\sqrt{2}}{16\pi} \int \frac{d\bar{\tau}}{d\bar{a}} \eta \chi \wedge (D_+ + 4\pi\lambda_+) + \frac{i\sqrt{2}}{2^7\pi} \int \frac{d\tau}{da} (\psi \wedge \psi) \wedge (4\pi\lambda_- + D_+) \right. \\
& \left. + \frac{1}{3 \cdot 2^{11}\pi i} \int \frac{d^2\tau}{da^2} \psi \wedge \psi \wedge \psi \wedge \psi \right],
\end{aligned} \tag{2.4}$$

where $\int_{Pic(X)}$ denotes a sum over line bundles and an integration over $\mathbf{T}^{b_1}$. The integration over ψ is understood as integration of differential forms on $\mathbf{T}^{b_1}$. The orientation of the measure of the finite-dimensional integral (2.4) corresponds to a choice of Donaldson orientation of the moduli space of instantons [15].

Consider now the generating function for an arbitrary insertion of zero, one, two and three observables. We introduce the formal sums of cycles

$$\gamma = \sum_{i=1}^{b_1} \zeta_i \delta_i, \quad S = \sum_{i=1}^{b_2} \lambda_i S_i, \quad \Sigma^3 = \sum_{i=1}^{b_3} \theta_i \Sigma_i^3, \tag{2.5}$$

where δ_i , $i = 1, \dots, b_1$, S_i , $i = 1, \dots, b_2$ and Σ_i^3 , $i = 1, \dots, b_3 = b_1$ are a basis of one, two, and three cycles, respectively. The basis of one-cycles δ_i is dual to $\beta_i \in H^1(X, \mathbb{Z})$. The λ_i are complex numbers, and ζ_i, θ_i are Grassmann variables. The insertion of observables corresponding to these cycles is

$$a_1 \int_\gamma Ku + a_2 \int_S K^2 u + a_3 \int_{\Sigma^3} K^3 u, \tag{2.6}$$

where a_1, a_2 and a_3 are constants that should be fixed by comparison to known mathematical results. The constant for the two-observable has been already fixed in [9], $a_2 = i/\sqrt{2}\pi$. K is the canonical descent operator in the normalization of [9], and we have, explicitly:

$$\begin{aligned}
Ku &= \frac{1}{4\sqrt{2}} \frac{du}{da} \psi, \\
K^2 u &= \frac{1}{32} \frac{d^2 u}{da^2} \psi \wedge \psi - \frac{\sqrt{2}}{4} \frac{du}{da} (F_+ + D), \\
K^3 u &= \frac{1}{\sqrt{2}2^7} \frac{d^3 u}{da^3} \psi \wedge \psi \wedge \psi - \frac{3}{16} \frac{d^2 u}{da^2} \psi \wedge (F_+ + D) - \frac{3\sqrt{2}i}{8} \frac{du}{da} (2d\chi - *d\eta).
\end{aligned} \tag{2.7}$$

In addition, we have to take into account various contact terms associated to the intersecting cycles [9][10]. These come from intersections of two, three and four cycles on the manifold X . For the intersection of two cycles we have

$$\begin{aligned} S \cap S &\in H_0(X, \mathbb{Z}), \quad \Sigma^3 \cap \gamma \in H_0(X, \mathbb{Z}), \\ \Sigma^3 \cap S &\in H_1(X, \mathbb{Z}), \quad \Sigma^3 \cap \Sigma^3 \in H_2(X, \mathbb{Z}). \end{aligned} \quad (2.8)$$

For intersection of three cycles, we have the possibilities

$$S \cap \Sigma^3 \cap \Sigma^3 \in H_0(X, \mathbb{Z}), \quad \Sigma^3 \cap \Sigma^3 \cap \Sigma^3 \in H_1(X, \mathbb{Z}), \quad (2.9)$$

and for intersection of four cycles we only have the possibility

$$\Sigma^3 \cap \Sigma^3 \cap \Sigma^3 \cap \Sigma^3 \in H_0(X, \mathbb{Z}). \quad (2.10)$$

The contact term corresponding to $S \cap S$ was obtained in [9]. A proposal for the structure of the contact terms corresponding to more general intersecting cycles was made in [10]. According to this proposal, the contact term associated to the intersection of p cycles is given by the appropriate descendant of the p -th derivative of the Seiberg-Witten prepotential with respect to a deformation parameter τ_0 . Moreover [11][12], this deformation parameter τ_0 can be related to the dynamically generated scale of the theory Λ (in the case of the asymptotically free theories) or to the microscopic gauge coupling, in the case of self-dual gauge theories. For $SU(2)$ $\mathcal{N} = 2$ supersymmetric Yang-Mills theory, the relation between τ_0 and Λ is given by $\Lambda^4 = e^{i\pi\tau_0}$. This is related to the fact that Λ can be identified with the first slow time of the Toda-Whitham hierarchy [16]. Following the proposal of [10] the contact terms for the various intersections in (2.8), (2.9) and (2.10) are of the form

$$\begin{aligned} &S^2 T(u) + a_{13} T(u) (\Sigma^3 \cap \gamma) + a_{32} \int_{\Sigma^3 \cap S} K T(u) + a_{33} \int_{\Sigma^3 \cap \Sigma^3} K^2 T(u) \\ &+ a_{332} \mathcal{F}_{\tau_0}^{(3)}(S \cap \Sigma^3 \cap \Sigma^3) + a_{333} \int_{\Sigma^3 \cap \Sigma^3 \cap \Sigma^3} K \mathcal{F}_{\tau_0}^{(3)} \\ &+ a_{3333} \mathcal{F}_{\tau_0}^{(4)}(\Sigma^3 \cap \Sigma^3 \cap \Sigma^3 \cap \Sigma^3), \end{aligned} \quad (2.11)$$

where the a 's are constants which will be determined below. In this equation, $\mathcal{F}_{\tau_0}^{(p)}$ denotes the p -th derivative of the prepotential with respect to τ_0 , and $T(u) = (4/\pi i) \mathcal{F}_{\tau_0}^{(2)}$. The contact term for $S \cap S$ can be written as [10][11]:

$$T(u) = \frac{1}{4} \left[2u - a \frac{du}{da} \right]. \quad (2.12)$$

The constants a_{ij} , a_{ijk} and a_{ijkl} will be obtained in terms of a_i , $i = 1, 2, 3$, using single-valuedness of the integrand on the u -plane. Notice that one expects, on physical grounds, that a_{ij} is proportional to $a_i a_j$, and so on.

We can already plug the observables (2.6) and the corresponding contact terms (2.11) into the generating function and write an explicit expression for the u -plane integral. It is important, however, to check that the resulting expression has good properties under duality transformations, that is, that the integrand is single-valued in the u -plane. This is not obvious due to the holomorphic “functions” that appear in (2.7) and (2.11). Following the strategy of [9], we will first of all integrate the auxiliary field D . Comparing to the expressions in the simply-connected case, we see that the new terms coupling to D appear in the combination

$$-\frac{i}{4\pi} \frac{du}{da} \left[(4\pi\lambda_- + D) \wedge \tilde{S} \right] \quad (2.13)$$

where

$$\tilde{S} = S - \frac{\sqrt{2}}{32} \frac{d\tau}{du} \psi \wedge \psi + \frac{3\pi}{4i} a_3 \frac{d^2 u}{da^2} \frac{da}{du} \psi \wedge \Sigma^3 - \sqrt{2}\pi i a_{33} \frac{dT}{du} \Sigma^3 \wedge \Sigma^3, \quad (2.14)$$

and we interpret Σ^3 as a harmonic one-form using Poincaré duality and the Hodge theorem. To guarantee that the resulting lattice sum over first Chern classes is well-behaved under duality transformations, the two-form $\tilde{S}$ should be invariant under duality. To achieve this, we redefine the ψ field as

$$\tilde{\psi} = \psi - \frac{12\pi}{\sqrt{2}i} a_3 \frac{d^2 u}{da^2} \frac{da}{d\tau} \Sigma^3. \quad (2.15)$$

Then, if we choose

$$a_{33} = -9\pi^2 a_3^2 \quad (2.16)$$

$\tilde{S}$ becomes

$$\tilde{S} = S - \frac{\sqrt{2}}{32} \frac{d\tau}{du} \tilde{\psi} \wedge \tilde{\psi} + \frac{9\pi^3 \sqrt{2}i}{4} a_3^2 \Sigma^3 \wedge \Sigma^3. \quad (2.17)$$

We then see that, if $\tilde{\psi}$ is a modular form of weight $(1, 0)$, then $\tilde{S}$ is a modular form of weight zero. Notice that the redefinition of ψ in (2.15) does not change the ψ measure. To obtain the above expression for $\tilde{S}$, we have taken into account that

$$4\pi i \frac{dT}{da} - \left(\frac{d^2 u}{da^2} \right)^2 \frac{da}{d\tau} = \pi i \frac{du}{da}. \quad (2.18)$$

Once this redefinition has been made, the u -plane integral involves a lattice sum identical to the one in the simply-connected case but with the substitution $S \rightarrow \tilde{S}$, and

additional holomorphic insertions (coming from the observables and contact terms) that, once they are reexpressed in terms of $\tilde{\psi}$ and $\tilde{S}$, should be modular forms of weight zero. This is in fact the case for appropriate choices of the constants in (2.11). The computation is lengthy but straightforward. Duality invariance fixes all the constants in (2.11) in terms of a_1 , a_2 and a_3 . The final result is:

$$\begin{aligned} a_{13} &= -6\pi^2 a_1 a_3, & a_{32} &= -6\sqrt{2}\pi i a_3, \\ a_{332} &= -72\sqrt{2}\pi i a_3^2, & a_{333} &= 36\pi^2 i a_3^3, \\ a_{3333} &= -216\pi^3 i a_3^4. \end{aligned} \quad (2.19)$$

The u -plane integral is therefore given by:

$$\begin{aligned} Z_u &= \int_{\Gamma^0(4) \backslash \mathcal{H}} \frac{dx dy}{y^{1/2}} \mu(\tau) \int_{Pic(X)} d\psi \exp \left[2pu + \tilde{S}^2 \hat{T}(u) + \frac{\sqrt{2}}{32} u \frac{d\tau}{du} (\tilde{S}, \tilde{\psi}^2) \right. \\ &\quad + \frac{a_1}{4\sqrt{2}} \frac{du}{da} \int_X \tilde{\psi} \wedge [\gamma] - 3\pi^2 a_1 a_3 u (\Sigma^3 \cap \gamma) - \frac{3\pi i}{8} a_3 \frac{du}{da} (\tilde{S}, \tilde{\psi} \wedge \Sigma^3) \\ &\quad + \frac{3\sqrt{2}\pi^3 i}{2} a_3^2 u (\tilde{S}, \Sigma^3 \wedge \Sigma^3) + \frac{7}{3 \cdot 2^{10}} u \left(\frac{d\tau}{du} \right)^2 \tilde{\psi}^4 - \frac{\sqrt{2}\pi i}{64} a_3 \frac{d\tau}{da} (\tilde{\psi}^3 \wedge \Sigma^3) \\ &\quad + \frac{9\pi^3 i}{64} a_3^2 u \frac{d\tau}{du} (\tilde{\psi}^2, \Sigma^3 \wedge \Sigma^3) - \frac{9\sqrt{2}\pi^4}{256} a_3^3 \frac{du}{da} \int_X \tilde{\psi} \wedge \Sigma^3 \wedge \Sigma^3 \wedge \Sigma^3 \\ &\quad \left. + \frac{135\pi^6}{16} a_3^4 u (\Sigma^3 \cap \Sigma^3 \cap \Sigma^3 \cap \Sigma^3) \right] \Psi(\tilde{S}) \end{aligned} \quad (2.20)$$

where

$$\begin{aligned} \mu(\tau) &= -\frac{\sqrt{2}}{2} \frac{da}{d\tau} A^\chi B^\sigma \\ \Psi(\tilde{S}) &= \exp(2i\pi\lambda_0^2) \exp \left[-\frac{1}{8\pi y} \left(\frac{du}{da} \right)^2 \tilde{S}_-^2 \right] \\ &\quad \sum_{\lambda \in H^2 + \frac{1}{2} w_2(E)} \exp \left[-i\pi\bar{\tau}(\lambda_+)^2 - i\pi\tau(\lambda_-)^2 + \pi i(\lambda - \lambda_0, w_2(X)) \right] \\ &\quad \exp \left[-i \frac{du}{da} (\tilde{S}_-, \lambda_-) \right] \left[(\lambda_+, \omega) + \frac{i}{4\pi y} \frac{du}{da} (\tilde{S}_+, \omega) \right] \\ \hat{T}(u) &= T(u) + \frac{1}{8\pi \text{Im}\tau} \left(\frac{du}{da} \right)^2. \end{aligned} \quad (2.21)$$

Notice that, in the above exponential, all the terms are modular forms of weight zero if $\tilde{\psi}$ is a modular form of weight $(1, 0)$. There is another check one can do of the above functions: one can formally assign an R -charge to the cycles in such a way that the observables have

R -charge zero, namely: $R(\gamma) = -3$, $R(S) = -2$, $R(\Sigma^3) = -1$. Taking into account that $R(\psi) = 1$, $R(a) = 2$, we see that the definitions of $\tilde{\psi}$, $\tilde{S}$ are consistent with this R -charge assignment and that, moreover, all the terms in the exponential of (2.20) have zero R -charge. This is in fact an example of Seiberg's trick since the insertions of (2.6)(2.11) may be regarded as operators in some UV theory (*e.g.* in a brane configuration).

The remaining constants a_1, a_3 will be fixed below by comparing to certain topological results [17]. In principle, the coefficients in (2.20) are fixed by the proposal of equation (2.14) of the first paper in [10]. Although we find the proposal of Losev, Nekrasov, and Shatashvili natural, because of the many conventions involved we have not checked that the coefficients derived in (2.20) are consistent with their equation (2.14).

3. Donaldson wall-crossing

In this section we want to compute the wall-crossing formulae associated to the region at infinity of the u -plane. To compare to mathematical results, it is better to write the expression in terms of ψ , S . There are two standard mathematical facts for manifolds of $b_2^+ = 1$ that will be very useful in writing the resulting wall-crossing formula [17]. First, for any $\beta_1, \beta_2, \beta_3$ and β_4 in $H^1(X, \mathbb{Z})$, one has $\beta_1 \wedge \beta_2 \wedge \beta_3 \wedge \beta_4 = 0$. Second, the image of the map

$$\wedge : H^1(X, \mathbb{Z}) \otimes H^1(X, \mathbb{Z}) \longrightarrow H^2(X, \mathbb{Z}) \quad (3.1)$$

is generated by a single rational cohomology class Σ (not to be confused with the three-cycle Σ^3). We introduce now the antisymmetric matrix a_{ij} associated to the basis β_i of $H^1(X, \mathbb{Z})$, $i = 1, \dots, b_1$, as $\beta_i \wedge \beta_j = a_{ij} \Sigma$. Finally, we introduce the two-form on $\mathbf{T}^{b_1}$ as

$$\Omega = \sum_{i < j} a_{ij} \beta_i^\# \wedge \beta_j^\#, \quad (3.2)$$

which does not depend on the choice of basis. This is a volume element for the torus, hence

$$\text{vol}(\mathbf{T}^{b_1}) = \int_{\mathbf{T}^{b_1}} \frac{\Omega^{b_1/2}}{(b_1/2)!}. \quad (3.3)$$

(For the manifolds under discussion b_1 is even.) As a simple example, if $X = \mathbb{CP}^1 \times F_g$, where F_g is a Riemann surface of genus g , then $\Sigma = [\mathbb{CP}^1]$ (the Poincaré dual to the two-homology class of $\mathbb{CP}^1$) and $\text{vol}(\mathbf{T}^{b_1}) = 1$.

With all these ingredients, we can already write the wall-crossing formulae. Notice that, because of the first fact, many terms on the u -plane integral (2.21) vanish. We

will write the formula first for an insertion of two and four observables. In this case, a straightforward generalization of the arguments in [9] gives:

$$\begin{aligned}
WC(\lambda) = & -32i(-1)^{(\lambda-\lambda_0, w_2(X))} e^{2\pi i \lambda_0^2} (8i)^{-b_1/2} \\
& \cdot \left[q^{-\lambda^2/2} \frac{da}{d\tau} \left((u^2 - 1) \frac{d\tau}{du} \right)^{1-\frac{b_1}{2}} \left(\frac{(\frac{2i}{\pi} \frac{du}{d\tau})^2}{u^2 - 1} \right)^{\sigma/8} \exp \left\{ 2pu + S^2 T(u) - i(\lambda, S)/h \right\} \right. \\
& \left. \int_{\mathbf{T}^{b_1}} d\psi \exp \left\{ \frac{i\sqrt{2}}{32} \left(\frac{d^2 u}{da^2} \frac{S}{2\pi} + \frac{d\tau}{da} \lambda, \psi^2 \right) \right\} \right]_{q^0}, \tag{3.4}
\end{aligned}$$

where we use the value obtained in [9] for the universal constants α , β , and

$$h(\tau) = \frac{da}{du} = \frac{1}{2} \vartheta_2 \vartheta_3. \tag{3.5}$$

We can actually compute the integral over $\mathbf{T}^{b_1}$ and give an expression in terms of modular forms which generalizes the expressions given in [9][18][19]. Using the explicit expressions given in [9] for the Seiberg-Witten solution in terms of modular forms, we find that

$$\frac{d^2 u}{da^2} = 4f_1(q), \quad \frac{d\tau}{da} = \frac{16i}{\pi} f_2(q), \tag{3.6}$$

where

$$\begin{aligned}
f_1(q) &= \frac{2E_2 + \vartheta_2^4 + \vartheta_3^4}{3\vartheta_4^8} = 1 + 24q^{1/2} + \dots, \\
f_2(q) &= \frac{\vartheta_2 \vartheta_3}{2\vartheta_4^8} = q^{1/8} + 18q^{5/8} + \dots.
\end{aligned} \tag{3.7}$$

Using (2.3) and (3.2) we find

$$\psi^2 = 2(\Sigma \otimes \Omega), \tag{3.8}$$

hence we can perform the integral over $\mathbf{T}^{b_1}$ to write a very explicit expression for (3.4):

$$\begin{aligned}
WC(\lambda) = & -\frac{i}{2}(-1)^{(\lambda-\lambda_0, w_2(X))} e^{2\pi i \lambda_0^2} 2^{-b_1/4} \text{vol}(\mathbf{T}^{b_1}) \\
& \cdot \left[q^{-\lambda^2/2} h(\tau)^{b_1-2} \vartheta_4^\sigma \exp \left\{ 2pu + S^2 T(u) - i(\lambda, S)/h \right\} \right. \\
& \left. \sum_{b=0}^{b_1/2} \frac{1}{(8i)^b} \binom{b_1/2}{b} f_1(q)^b f_2(q)^{b_1/2-b-1} (S, \Sigma)^b (\lambda, \Sigma)^{b_1/2-b} \right]_{q^0}. \tag{3.9}
\end{aligned}$$

Notice that this result confirms the conjecture in p. 18 of [17]. As a check of this expression, and also to fix an overall coefficient depending on b_1 , we will compute the wall-crossing for

the correlator $p^r S^{d-2r}$ on a ruled surface, where d is half the dimension of the moduli space, and compare it to the expressions in [17]. Introduce the integer cohomology class $\zeta = 2\lambda$. In [17], Muñoz computes the wall-crossing for walls satisfying $\zeta^2 = p_1$ and $\zeta^2 = p_1 + 4$, where p_1 is the Pontryagin number of the instanton bundle. For $\zeta^2 = p_1$, it is easy to see that only the first coefficients in the expansion in q contribute to (3.9), and there is no contribution from the contact term $T(u)$. To compare the formulae, notice that we have to multiply our wall-crossing expression by $r!(d-2r)!$, as we are considering the Donaldson-Witten generating function. We finally obtain,

$$\delta_{\zeta, \lambda_0}(p^r S^{d-2r}) = \frac{1}{2}(-1)^{(\lambda - \lambda_0, w_2(X))}(-i)^{b_1/2} 2^{-3b_1/4 - b - d} (-1)^{r+d} p^r \text{vol}(\mathbf{T}^{b_1}) \sum_{b=0}^{b_1/2} \frac{(b_1/2)!}{(b_1/2 - b)!} \binom{d-2r}{b} (S, \zeta)^{d-2r-b} (S, \zeta)^b (\zeta, \Sigma)^{b_1/2-b}. \quad (3.10)$$

If we compare with [17], we find perfect agreement except for an overall factor $1/2$ (a standard discrepancy between topological and quantum field theory normalizations), and a factor $(-i)^{b_1/2} 2^{-3b_1/4}$. The latter factor is due to the normalization of the fermions in the physical theory. In order to make the identification in (2.3) and to use the normalization of topologists, we have to correct the ψ measure with an overall factor $i^{b_1/2} 2^{9b_1/4}$. As we will see in the next section, with this normalization the above formula gives the right answer for the generalized Seiberg-Witten wall-crossing formula of [20][21]. The case $\zeta^2 = p_1 + 4$ also agrees with [17]. In this case the computation is more involved, as one has to take into account the first two coefficients in the q -expansion of the various functions in (3.9).

We consider now an arbitrary insertion of observables associated to one, two, and three cycles. We have new contact terms as well as new terms in the integration over the torus. To write these in a convenient way, notice that we can use Poincaré duality and the isomorphism $H_1(X, \mathbb{Z}) \simeq H^1(\mathbf{T}^{b_1}, \mathbb{Z})$ to obtain a one-form $\Sigma^{3\sharp}$ in $H^1(\mathbf{T}^{b_1}, \mathbb{Z})$. Define

$$\iota_{\beta_k^\sharp} \Omega = \sum_p a_{kp} \beta_p^\sharp. \quad (3.11)$$

Using (2.3), we find

$$\psi \wedge \Sigma^3 = (\iota_{\Sigma^{3\sharp}} \Omega) \Sigma, \quad (3.12)$$

In the same way, using the isomorphism $H_1(X, \mathbb{Z}) \simeq H^1(X, \mathbb{Z})$ given by $\delta_i \rightarrow \beta_i$, we can define $\gamma^\sharp = \sum_{i=1}^{b_1} \zeta_i \beta_i^\sharp$, where the ζ_i were defined in (2.5). We then obtain, using (2.3) again,

$$\int_\gamma \psi = \gamma^\sharp. \quad (3.13)$$

The functions appearing in the u -plane integral can be written in terms of Jacobi theta functions and Eisenstein series as follows,

$$\frac{dT}{da} = -8f_3(q), \quad \mathcal{F}_{\tau_0}^{(3)} = -\frac{\pi^2}{4}f_4(q), \quad (3.14)$$

where

$$\begin{aligned} f_3(q) &= \frac{1}{16\vartheta_2\vartheta_3} \left[\frac{1}{9\vartheta_4^8} (2E_2 + \vartheta_2^4 + \vartheta_3^4)^2 - 1 \right] = q^{3/8} + 12q^{7/8} + \dots, \\ f_4(q) &= \frac{1}{8(\vartheta_2\vartheta_3)^2} \left[\frac{1}{2}(\vartheta_2^4 + \vartheta_3^4) - E_2 + \frac{1}{54\vartheta_4^8} (2E_2 + \vartheta_2^4 + \vartheta_3^4)^3 \right] = q^{1/4} + 8q^{3/4} + \dots. \end{aligned} \quad (3.15)$$

The wall-crossing formula now reads,

$$\begin{aligned} WC(\lambda) &= -\frac{i}{2}(-1)^{(\lambda-\lambda_0, w_2(X))} e^{2\pi i \lambda_0^2} 2^{7b_1/4} (-i\pi)^{b_1/2} \\ &\cdot \left[q^{-\lambda^2/2} h(\tau)^{b_1-2} f_2(q)^{-1} \vartheta_4^\sigma \exp \left\{ 2pu + (S^2 - 6\pi^2 a_1 a_3 (\Sigma^3 \cap \gamma)) T(u) \right. \right. \\ &\quad \left. \left. + 18\sqrt{2}\pi^3 i a_3^2 (\Sigma^3 \cap \Sigma^3 \cap S) f_4(q) - i(\lambda, S)/h - 72\sqrt{2}\pi^3 a_3^2 f_3(q) (\Sigma^3 \wedge \Sigma^3, \lambda) \right\} \right. \\ &\quad \left. \int_{\mathbf{T}^{b_1}} \exp \left[2(P(q), \Sigma) \Omega + (R(q), \Sigma) \iota_{\Sigma^3 \#} \Omega + Q(q) \gamma^\# \right] \right]_{q^0}, \end{aligned} \quad (3.16)$$

where

$$\begin{aligned} P(q) &= \frac{i\sqrt{2}}{16\pi} (f_1(q)S + 8if_2(q)\lambda), \\ R(q) &= 3\pi a_3 (4if_3(q)S - f_1(q)\lambda), \\ Q(q) &= \frac{\sqrt{2}a_1}{8} \frac{1}{h}. \end{aligned} \quad (3.17)$$

In the case of ruled surfaces and for $\zeta^2 = p_1$, we find again agreement with the expressions obtained by Muñoz. In fact, comparing to the formula in p.13 of [17], we can find the values of a_1, a_3 :

$$a_1 = \pi^{-1/2} 2^{3/4} e^{-\frac{\pi i}{4}}, \quad a_3 = \frac{\pi^{-3/2}}{6} 2^{1/4} e^{\frac{\pi i}{4}}. \quad (3.18)$$

4. The Seiberg-Witten contribution

We can now follow the strategy in [9] and find the form of the Seiberg-Witten contribution at the monopole and dyon cusps. We focus on the monopole cusp, as the contribution

at the dyon cusp can be obtained using the $\mathbb{Z}_2$ symmetry on the u -plane. In fact, as the functions involved in the monopole contribution are universal and they have been obtained in [9] in the simply-connected case, we will be able to *derive* the general wall-crossing formula of [20][21].

A crucial ingredient in the discussion of the Seiberg-Witten contribution for non-simply connected manifolds is that we have to consider generalized Seiberg-Witten invariants, *i.e.*, we have to consider correlation functions of one-observables. Recall that the basic observable in Seiberg-Witten theory is the two-form a_D on $\mathcal{M}_\lambda$. The first descendant of a_D (in the topological field theory associated to the Seiberg-Witten monopole equations) is ψ , which is a one-form on X and also a one-form on $\mathcal{M}_\lambda$. It can be written as:

$$\psi = \sum_{i=1}^{b_1} \nu_i \beta_i, \quad (4.1)$$

where $\beta_i \in H^1(X, \mathbb{Z})$, $i = 1, \dots, b_1$ is the basis of one-forms considered before, and

$$\nu_i = \int_{\delta_i} \psi \quad (4.2)$$

are the one-observables of Seiberg-Witten theory. The generalized Seiberg-Witten invariant associated to the one-forms $\beta_1, \dots, \beta_r$ is defined by intersection theory on $\mathcal{M}_\lambda$:

$$SW(\lambda, \beta_1 \wedge \dots \wedge \beta_r) = \int_{\mathcal{M}_\lambda} \nu_1 \wedge \dots \wedge \nu_r \wedge a_D^{\frac{d_\lambda - r}{2}}, \quad (4.3)$$

where $d_\lambda = \lambda^2 - (2\chi + 3\sigma)/4$ is the virtual dimension of $\mathcal{M}_\lambda$. These generalized invariants (and their wall-crossing) have been considered in [21].

The Seiberg-Witten twisted Lagrangian near $u = 1$ (with the monopole fields included) can be written as [9][11]

$$\begin{aligned} & \{\bar{\mathcal{Q}}, W\} + \frac{i}{16\pi} \tilde{\tau}_M F \wedge F + p(u) \text{Tr} R \wedge R \\ & + \ell(u) \text{Tr} R \wedge \tilde{R} - \frac{i\sqrt{2}}{2^7 \cdot \pi} \frac{d\tilde{\tau}_M}{da_D} (\psi \wedge \psi) \wedge F \\ & + \frac{i}{3 \cdot 2^{11} \pi} \frac{d^2 \tilde{\tau}_M}{da_D^2} \psi \wedge \psi \wedge \psi \wedge \psi, \end{aligned} \quad (4.4)$$

and, as we see, the terms which are not $\mathcal{Q}$ -exact do not depend on the metric. The exponentiation of the terms involving the densities $\text{Tr} R \wedge R$, $\text{Tr} R \wedge \tilde{R}$ gives, after integration on X , the gravitational factors $P(u)^{\sigma/8}$, $L(u)^{\chi/4}$ considered in [9]. The term involving $F \wedge F$

gives a factor $C(u)^{\lambda^2/2}$, where $F = 4\pi\lambda$ and $C(u) = e^{-2\pi i\tilde{\tau}_M}$. The terms $C(u)$, $L(u)$ and $P(u)$ are universal (they do not depend on the manifold X) and they were found in [9] using matching of wall-crossing in the simply-connected case. They can be written as

$$\begin{aligned} L(u) &= \pi i \alpha^4 \left((u^2 - 1) \frac{d\tau_D}{du} \right), \\ P(u) &= -\pi^2 \beta^8 a_D^{-1} (u^2 - 1), \\ C(u) &= \frac{a_D}{q_D}, \end{aligned} \tag{4.5}$$

where $q_D = e^{2\pi i\tau_D}$. The last relation tells us that the gauge coupling $\tilde{\tau}_M$ appearing in (4.4) is given by

$$\tilde{\tau}_M = \tau_D - \frac{1}{2\pi i} \log a_D, \tag{4.6}$$

and therefore it is smooth at the monopole cusp. This defines the prepotential $\tilde{\mathcal{F}}_M(a_D)$ through the equation $\tilde{\mathcal{F}}_M''(a_D) = \tilde{\tau}_M$.

First we analyze the Seiberg-Witten contribution when only two and four observables are introduced. It can be written as

$$\begin{aligned} \langle e^{p\mathcal{O}+I_2(S)} \rangle_{\lambda, u=1} &= \int_{\mathcal{M}_\lambda} 2e^{2i\pi(\lambda_0 \cdot \lambda + \lambda_0^2)} C(u)^{\lambda^2/2} P(u)^{\sigma/8} L(u)^{\chi/4} \\ &\cdot \exp \left(2pu + i \frac{du}{da_D} (S, \lambda) + S^2 T_M(u) \right) \exp \left[(P_M(u), \Sigma) \sum_{i,j=1}^{b_1} a_{ij} \nu_i \nu_j \right], \end{aligned} \tag{4.7}$$

where

$$P_M(u) = \frac{i\sqrt{2}}{32} \left(\frac{d^2 u}{da_D^2} \frac{S}{2\pi} + \frac{d\tilde{\tau}_M}{da_D} \lambda \right), \tag{4.8}$$

which is the magnetic version of the function in the $\mathbf{T}^{b_1}$ integral of (3.4), but with the smooth coupling constant (4.6). When we expand the exponential involving the one-observables, we obtain the generalized Seiberg-Witten invariants with $2b$ insertions, $b = 0, \dots, b_1$. The final expression is:

$$\begin{aligned} \langle e^{p\mathcal{O}+I_2(S)} \rangle_{\lambda, u=1} &= \sum_{b=0}^{b_1/2} \frac{1}{b!} \text{Res}_{a_D=0} \left[2e^{2i\pi(\lambda_0 \cdot \lambda + \lambda_0^2)} C(u)^{\lambda^2/2} P(u)^{\sigma/8} L(u)^{\chi/4} \right. \\ &\cdot \exp \left(2pu + i \frac{du}{da_D} (S, \lambda) + S^2 T_M(u) \right) a_D^{-d_\lambda/2+b-1} (P_M(u), \Sigma)^b \Big] \\ &\cdot \sum_{i_p, j_p=1}^{b_1} a_{i_1 j_1} \cdots a_{i_b j_b} SW(\lambda, \beta_{i_1} \wedge \beta_{j_1} \wedge \cdots \wedge \beta_{i_b} \wedge \beta_{j_b}). \end{aligned} \tag{4.9}$$

On the other hand, the wall-crossing formula for the u -plane integral near $u = 1$ can be written as

$$\begin{aligned}
WC(\lambda) = & 2\pi i 2^{11b_1/4} i^{b_1/2} \alpha^\chi \beta^\sigma e^{2i\pi(\lambda_0 \cdot \lambda + \lambda_0^2)} \text{vol}(\mathbf{T}^{b_1}) \\
& \cdot \text{Res}_{a_D=0} \left[q_D^{-\lambda^2/2} \left((u^2 - 1) \frac{d\tau}{du} \right)^{\chi/4} (u^2 - 1)^{\sigma/8} \right. \\
& \cdot \left. \exp \left(2pu + i \frac{du}{da_D} (S, \lambda) + S^2 T_M(u) \right) (P(q_D), \Sigma)^{b_1/2} \right],
\end{aligned} \tag{4.10}$$

where we have included the extra factors depending on b_1 that we obtained in the previous section by comparing to the expressions in [17]. Notice that, from (4.6), one has

$$P(q_D) = P_M(u) + \frac{\sqrt{2}}{64\pi} \frac{1}{a_D} \lambda, \tag{4.11}$$

and (4.10) can then be expanded as:

$$\begin{aligned}
WC(\lambda) = & 2\pi i \left(\frac{i}{\pi} \right)^{b_1/2} \alpha^\chi \beta^\sigma e^{2i\pi(\lambda_0 \cdot \lambda + \lambda_0^2)} \text{vol}(\mathbf{T}^{b_1}) \\
& \cdot \sum_{b=0}^{b_1/2} \binom{b_1/2}{b} \text{Res}_{a_D=0} \left[q_D^{-\lambda^2/2} \left((u^2 - 1) \frac{d\tau}{du} \right)^{\chi/4} (u^2 - 1)^{\sigma/8} \right. \\
& \cdot \exp \left(2pu + i \frac{du}{da_D} (S, \lambda) + S^2 T_M(u) \right) \\
& \cdot \left. 2^{11b/2} \pi^b (P_M(u), \Sigma)^b (\lambda, \Sigma)^{b_1/2-b} a_D^{b-b_1/2} \right].
\end{aligned} \tag{4.12}$$

Now we can compare the expressions for wall-crossing obtained from (4.9) and (4.12). Notice again that, to identify the fields ψ with the one-observables in (4.2) we have to be careful with possible normalization factors needed to agree with the normalization used by topologists. But for the terms with no insertions, *i.e.* $b = 0$ in (4.9), we should be able to obtain the wall-crossing formula for this Seiberg-Witten invariant. In fact, we find:

$$WC(SW(\lambda)) = (-1)^{b_1/2} (\lambda, \Sigma)^{b_1/2} \text{vol}(\mathbf{T}^{b_1}). \tag{4.13}$$

As $\lambda \in H^2(X, \mathbb{Z}) + \frac{1}{2}w_2(X)$, and 2λ is the determinant line bundle of the corresponding Spin^c structure, we find perfect agreement with [20][21] (notice that the wall-crossing formula in Theorem 1.2 of [20] should have an extra factor of $2^{-b_1/2}$). For the wall-crossing of Seiberg-Witten invariants with one-observable insertions, the general formula of [21] is (in our notation):

$$WC(SW(\lambda, \beta_1 \wedge \cdots \wedge \beta_r)) = \frac{(-1)^{\frac{b_1-r}{2}}}{\left(\frac{b_1-r}{2}\right)!} (\lambda, \Sigma)^{\frac{b_1-r}{2}} \int_{\mathbf{T}^{b_1}} \beta_1^\# \wedge \cdots \wedge \beta_r^\# \wedge \Omega^{\frac{b_1-r}{2}}, \tag{4.14}$$

and we see that, if we introduce a normalization factor $2^{-9/4}\pi^{-1/2}i$ for the fields ψ , we obtain from the matching of wall-crossing the expression

$$\sum_{i_p, j_p=1}^b a_{i_1 j_1} \cdots a_{i_b j_b} WC(SW(\lambda, \beta_{i_1} \wedge \beta_{j_1} \wedge \cdots \wedge \beta_{i_b} \wedge \beta_{j_b})) =$$

$$2^b \frac{(-1)^{b_1/2-b}}{(b_1/2-b)!} (\lambda, \Sigma)^{b_1/2-b} \text{vol}(\mathbf{T}^{b_1}), \quad (4.15)$$

again in agreement with (4.14). The expression (4.14) can be actually derived by considering general insertions of one and three-observables.

We then see that, in general, the Donaldson invariants for a non-simply connected manifold should be written in terms of the generalized Seiberg-Witten invariants, and only in this case we have consistent formulae for matching of wall-crossing. When we include arbitrary observables associated to one- and three-cycles, the above expressions are more complicated and we have to take into account the terms involving γ and Σ^3 , as well as the new contact terms.

For manifolds of $b_2^+ > 1$, all the contact terms appear in the Seiberg-Witten contribution. They are given by the expression in (2.11), where the prepotential is now the prepotential $\tilde{\mathcal{F}}_M(a_D)$ (notice that, as all the contact terms are regular at the monopole cusp, we can take as well the dual prepotential $\mathcal{F}_D(a_D)$). In general we have a complicated expression which can be written explicitly using the previous results. In the simple type case, $d_\lambda = 0$, the Seiberg-Witten moduli space consists of a finite set of points and counting them with appropriate signs we obtain $SW(\lambda)$. We only have to compute the different functions at $u = 1$ (the first term in the expansion in a_D). Using also (3.18), we can already write an explicit expression for the SW contribution of λ to the Donaldson invariants at $u = 1$, with an arbitrary insertion of observables:

$$\langle e^{p\mathcal{O}+I_2(S)+I_1(\gamma)+I_3(\Sigma^3)} \rangle_{\lambda, u=1} = 2^{1+\frac{7\chi}{4}+\frac{11\sigma}{4}} e^{2i\pi(\lambda_0 \cdot \lambda + \lambda_0^2)}$$

$$\cdot \exp\left\{2p + \frac{1}{2}S^2 - (\Sigma^3 \cap \gamma) - \frac{1}{4}(\Sigma^3 \wedge \Sigma^3, S) + \frac{1}{96}(\Sigma^3 \cap \Sigma^3 \cap \Sigma^3 \cap \Sigma^3)\right\} \quad (4.16)$$

$$\cdot \exp\left\{2(S, \lambda) - \frac{1}{4}(\Sigma^3 \wedge \Sigma^3, \lambda)\right\} SW(\lambda).$$

We see that the contact terms give, on the one hand, new contributions which depend on the intersection theory on X . On the other hand, there is a term coming from the intersection of the two-cycle $\Sigma^3 \cap \Sigma^3$ with the Poincaré dual of the basic class λ , as was

suspected in [6] using the cosmic string approach. Notice that all the coefficients in (4.16) are real, as needed for consistency.

We can now write an expression for the Donaldson invariants of non-simply connected manifolds of simple type, summing over the basic classes and considering both the monopole and the dyon contributions. Using the $\mathbf{Z}_2$ symmetry on the u -plane, and taking into account the R -charges of the different terms in (4.16), we obtain: ²

$$\begin{aligned}
& \langle e^{p\mathcal{O}+I_2(S)+I_1(\gamma)+I_3(\Sigma^3)} \rangle \\
&= 2^{1+\frac{7\chi}{4}+\frac{11\sigma}{4}} \left(\exp \left\{ 2p + \frac{1}{2}S^2 - (\Sigma^3 \cap \gamma) - \frac{1}{4}(\Sigma^3 \wedge \Sigma^3, S) + \frac{1}{96}(\Sigma^3 \cap \Sigma^3 \cap \Sigma^3 \cap \Sigma^3) \right\} \right. \\
&\quad \cdot \sum_{\lambda} e^{2i\pi(\lambda_0 \cdot \lambda + \lambda_0^2)} SW(\lambda) \exp \left\{ 2(S, \lambda) - \frac{1}{4}(\Sigma^3 \wedge \Sigma^3, \lambda) \right\} \\
&\quad + i^{\frac{\chi+\sigma}{4}-w_2^2(E)} \exp \left\{ -2p - \frac{1}{2}S^2 + (\Sigma^3 \cap \gamma) + \frac{1}{4}(\Sigma^3 \wedge \Sigma^3, S) - \frac{1}{96}(\Sigma^3 \cap \Sigma^3 \cap \Sigma^3 \cap \Sigma^3) \right\} \\
&\quad \cdot \sum_{\lambda} e^{2i\pi(\lambda_0 \cdot \lambda + \lambda_0^2)} SW(\lambda) \exp \left\{ -2i(S, \lambda) + \frac{i}{4}(\Sigma^3 \wedge \Sigma^3, \lambda) \right\} \Bigg).
\end{aligned} \tag{4.17}$$

5. Donaldson invariants for product ruled surfaces

In this section we will use the above results and the formulae in section 8 of [9] to write general expressions for the Donaldson invariants of product ruled surfaces $X = \mathbb{C}P^1 \times F_g$, where F_g is a Riemann surface of genus g . ³ For these manifolds, $b_1 = 2g$, $b_2 = 2$, $b_2^+ = 1$, so $\sigma = 0$ and $\chi = 4 - 2b_1$. $H^2(X, \mathbb{Z})$ is generated by the cohomology classes $[\mathbb{C}P^1]$, $[F_g]$, with intersection form $II^{1,1}$. Consider a general period point ω given by

$$\omega(\theta) = \frac{1}{\sqrt{2}}(e^{\theta}[\mathbb{C}P^1] + e^{-\theta}[F_g]). \tag{5.1}$$

The standard constant curvature metrics for $\mathbb{C}P^1, F_g$ give natural representatives for $[\mathbb{C}P^1], [F_g]$. Thus, choosing coordinates $z \in \mathbb{C}$ for $\mathbb{C}P^1$ and representing F_g (for $g > 1$) as a quotient by a Fuchsian group of the the Poincaré disk: $\mathcal{D} = \{w : |w| < 1\}$, we have

$$\begin{aligned}
[\mathbb{C}P^1] &= \frac{i}{2\pi(g-1)} \frac{dw \wedge d\bar{w}}{(1-|w|^2)^2}, \\
[F_g] &= \frac{i}{2\pi} \frac{dz \wedge d\bar{z}}{(1+|z|^2)^2},
\end{aligned} \tag{5.2}$$

² Curiously, the coefficient of $(\Sigma^3)^4$ is $1/96 = 2^{-5} \cdot 3^{-1}$. The factor of 3 would seem to imply divisibility properties by 3 for certain intersection numbers. These are easily confirmed in simple examples, but we did not find a general proof.

³ This case has also been discussed in eq. (2.15) of [10].

A metric with period point (5.1) then has scalar curvature $8\pi(e^\theta - e^{-\theta}(g-1))$ and will hence be positive for $e^{2\theta} > g-1$. By [7] there are no SW contributions to the Donaldson invariants for this metric. Thus, to compute the invariants in the chamber corresponding to the limit of a small volume for $\mathbb{C}P^1$ (with θ positive and large), we need only evaluate the u -plane integral.

The value of the integral depends on the non-abelian magnetic flux $w_2(E)$. The vanishing argument of section 5 of [9] applies when $w_2(E) \cdot [\mathbb{C}P^1] \neq 0$, (i.e. there is nontrivial flux through the small rational fiber). In this case we have simply that $Z_u = 0$. Now consider $w_2(E) = 0$. Here we can use the computation of section 8 of [9]. To do this, we choose the primitive null vector $z = [\mathbb{C}P^1] = \Sigma$. The expression derived in [9] will be valid as long as $z_+^2 = (z, \omega)^2$ is small, which is the case in the chamber under consideration (corresponding to θ large and positive). If we consider the expression (2.20) on a product ruled surface, we see that it involves a lattice sum $\Psi(\tilde{S})$ and a term involving the observables and the measure. From this last term we can derive an expression generalizing the holomorphic function f appearing in section 8 of [9]:

$$\begin{aligned} f = & \frac{\sqrt{2}}{8\pi i} (8i)^{1-g} \left((u^2 - 1) \frac{d\tau}{du} \right)^{1-g} \frac{du}{d\tau} \\ & \cdot \exp \left[2pu + \left(S^2 - \frac{1}{8} (S, \Sigma^3 \wedge \Sigma^3) \right) T(u) + \frac{\sqrt{2}}{64} a \frac{d\tau}{da} (S, \tilde{\psi}^2) \right. \\ & + \frac{a_1}{4\sqrt{2}} \frac{du}{da} \int_X \tilde{\psi} \wedge [\gamma] - u(\Sigma^3 \cap \gamma) - \frac{3\pi i}{8} a_3 \frac{du}{da} (S, \tilde{\psi} \wedge \Sigma^3) \\ & \left. - \frac{1}{12} u(S, \Sigma^3 \wedge \Sigma^3) \right], \end{aligned} \quad (5.3)$$

where we have taken into account the relation (2.17), and a_3 is given by (3.18). We also have that $(\tilde{S}, z) = (S, \Sigma)$. Using the computation leading to equation (8.15) of [9], we see that the Donaldson invariants of product ruled surfaces in the limit of a small volume for $\mathbb{C}P^1$ and for $w_2(E) = 0$ are given by

$$-8\sqrt{2}\pi i \left[\left(\int_{\mathbf{T}^{b_1}} d\tilde{\psi} f \right) h \coth \left(i \frac{(S, [\mathbb{C}P^1])}{2h} \right) \right]_{q^0} \quad (5.4)$$

When we consider only insertions of zero and two-observables, the integration over the torus is easily done and we obtain the following expression for the Donaldson invariants of $\mathbb{C}P^1 \times F_g$, which generalizes the expression obtained in [19][9] for $\mathbb{C}P^1 \times \mathbb{C}P^1$:

$$-16i \left[h(u^2 - 1) e^{2pu + S^2 T(u)} \left(2h^2 f_1(q)(S, [\mathbb{C}P^1]) \right)^g \coth \left(i \frac{(S, [\mathbb{C}P^1])}{2h} \right) \right]_{q^0}. \quad (5.5)$$

If $w_2(E) = [\mathbb{C}P^1]$, one can analyze the lattice reduction as in [9] including this nonzero flux, which gives extra phases in the lattice theta function. The effect of these phases is simply to change the $\coth\left(i(S, [\mathbb{C}P^1])/2h\right)$ in (5.4) by $-i \csc\left((S, [\mathbb{C}P^1])/2h\right)$, and we then obtain the generating function for $\mathbb{C}P^1 \times F_g$ and with $w_2(E) = [\mathbb{C}P^1]$:

$$-16 \left[h(u^2 - 1) e^{2pu + S^2 T(u)} \left(2h^2 f_1(q)(S, [\mathbb{C}P^1]) \right)^g \csc\left(\frac{(S, [\mathbb{C}P^1])}{2h}\right) \right]_{q^0}, \quad (5.6)$$

where we have included only zero and two-observables.

For applications to Floer theory and Gromov-Witten theory the most interesting chamber is on the other side of the Kähler cone, namely when $\theta \rightarrow -\infty$ giving a very small volume to F_g . We can obtain the Donaldson invariants for this chamber by summing over all the wall-crossing discontinuities. In general, denoting the lift $2\lambda_0$ of $w_2(E)$ by $\lambda_0 = \frac{1}{2}\epsilon_1[\mathbb{C}P^1] + \frac{1}{2}\epsilon_2[F_g]$ with $\epsilon_{1,2} = 0, 1$ we have walls at $\omega \cdot \lambda_{n,m} = 0$ where :

$$\lambda_{n,m} = (n + \frac{1}{2}\epsilon_1)[\mathbb{C}P^1] + (m + \frac{1}{2}\epsilon_2)[F_g] \quad (5.7)$$

for $(n + \frac{1}{2}\epsilon_1)(m + \frac{1}{2}\epsilon_2) < 0$, $n, m \in \mathbb{Z}$.⁴

The infinite sums over wall-crossings can be written explicitly in terms of modular forms using a result of Kronecker [22]:⁵

$$\begin{aligned} \sum_{m=1}^{\infty} \sum_{n=1}^{\infty} q^{mn} [e^{2\pi i(n\theta_1 + m\theta_2)} - e^{-2\pi i(n\theta_1 + m\theta_2)}] &= -1 + \frac{1}{1 - e^{-2\pi i\theta_1}} + \frac{1}{1 - e^{-2\pi i\theta_2}} \\ &\quad - i\eta^3(\tau) \frac{\vartheta_1(\theta_1 + \theta_2|\tau)}{\vartheta_1(\theta_1|\tau)\vartheta_1(\theta_2|\tau)}. \end{aligned} \quad (5.8)$$

We will consider in detail the cases $2\lambda_0 = [\mathbb{C}P^1]$, $2\lambda_0 = [\mathbb{C}P^1] + [F_g]$, which are relevant for the Floer cohomology of $F_g \times \mathbf{S}^1$ [2][24] as well as to the quantum cohomology of the moduli space of stable bundles over F_g [3][25]. We consider only insertions of zero, one, and two observables (*i.e.* we put $\Sigma^3 = 0$). As $w_2(X) = 0$ for product ruled surfaces, the only λ -dependent factors in the wall-crossing expression (3.16) are

$$q^{-\lambda^2/2} \exp\left[-\frac{i}{h}(\tilde{S}, \lambda)\right], \quad (5.9)$$

⁴ We require $\lambda^2 < 0$ rather than $\lambda^2 \leq 0$ because the “walls” at $\lambda^2 = 0$ are on the light-cone, and are never crossed when moving from one chamber to another.

⁵ Curiously, this identity showed up in recent studies of elliptic quantum cohomology, [23].

where $\tilde{S}$ has been defined in (2.14). Define now the formal variables

$$\begin{aligned} 2\pi\theta_1 &= \frac{1}{h}([\mathbb{C}P^1], \tilde{S}) = \frac{1}{h}([\mathbb{C}P^1], S), \\ 2\pi\theta_2 &= -\frac{1}{h}([F_g], \tilde{S}) = -\frac{1}{h}([F_g], S) + \frac{\sqrt{2}}{16} \frac{d\tau}{da} \Omega. \end{aligned} \quad (5.10)$$

The sum of wall-crossings can then be written, for $2\lambda_0 = [\mathbb{C}P^1]$, as

$$\begin{aligned} & - \sum_{m=1}^{\infty} \sum_{n=1}^{\infty} q^{m(n-1/2)} \left[e^{2\pi i((n-1/2)\theta_1 + m\theta_2)} - e^{-2\pi i((n-1/2)\theta_1 + m\theta_2)} \right] \\ &= \frac{i}{2 \sin(\pi\theta_1)} + i\eta^3(\tau) \frac{\vartheta_4(\theta_1 + \theta_2|\tau)}{\vartheta_1(\theta_1|\tau)\vartheta_4(\theta_2|\tau)}, \end{aligned} \quad (5.11)$$

and for $2\lambda_0 = [\mathbb{C}P^1] + [F_g]$ as

$$\begin{aligned} & - \sum_{m=1}^{\infty} \sum_{n=1}^{\infty} q^{(m-1/2)(n-1/2)} \left[e^{2\pi i((n-1/2)\theta_1 + (m-1/2)\theta_2)} - e^{-2\pi i((n-1/2)\theta_1 + (m-1/2)\theta_2)} \right] \\ &= i\eta^3(\tau) \frac{\vartheta_1(\theta_1 + \theta_2|\tau)}{\vartheta_4(\theta_1|\tau)\vartheta_4(\theta_2|\tau)}. \end{aligned} \quad (5.12)$$

These explicit expressions are obtained from (5.8) by shifting θ_1, θ_2 appropriately. The quotients of theta functions can be written in terms of Weierstrass σ functions, as in the blow-up formulae of [26]:

$$\begin{aligned} \frac{\vartheta_1(\theta|\tau)}{\vartheta_1'(0|\tau)} &= \frac{1}{\omega_2} e^{-\eta_2 \omega_2 \theta^2} \sigma(t), \\ \frac{\vartheta_4(\theta|\tau)}{\vartheta_4(0|\tau)} &= e^{-\eta_2 \omega_2 \theta^2} \sigma_3(t), \end{aligned} \quad (5.13)$$

where $t = \omega_2 \theta$ and ω_2 corresponds to the a -period of the Seiberg-Witten curve, $\omega_2 = \frac{8\pi}{\sqrt{2}} h$. The value of the Weierstrass zeta function at ω_2 can be written in terms of $E_2(\tau)$ as

$$\eta_2 = \frac{\pi^2}{6\omega_2} E_2(\tau). \quad (5.14)$$

In fact, the terms in (5.13) involving η_2 cancel the E_2 factors in the wall-crossing formula, and the resulting expressions are modular forms of weight zero (after integrating on $\mathbf{T}^{b_1}$).

The coefficients in the expansion of the σ functions only depend on the zero-observable u . After writing the Seiberg-Witten elliptic curve in Weierstrass form, one has that

$$g_2 = \frac{1}{4} \left(\frac{u^2}{3} - \frac{1}{4} \right), \quad g_3 = \frac{1}{48} \left(\frac{2u^3}{9} - \frac{u}{4} \right), \quad (5.15)$$

and the root relevant for $\sigma_3(t)$ is $e_3 = -u/12$. We then have the expansions

$$\begin{aligned}\sigma(t) &= t - \frac{g_2 t^5}{2^4 \cdot 3 \cdot 5} + \dots, \\ \sigma_3(t) &= 1 + \frac{u}{24} t^2 + \dots.\end{aligned}\tag{5.16}$$

Using (5.13) we find

$$\begin{aligned}\eta^3(\tau) \frac{\vartheta_4(\theta_1 + \theta_2|\tau)}{\vartheta_1(\theta_1|\tau)\vartheta_4(\theta_2|\tau)} &= -\frac{4}{\sqrt{2}} h e^{-2\eta_2 \omega_2 \theta_1 \theta_2} \frac{\sigma_3(t_1 + t_2)}{\sigma(t_1)\sigma_3(t_2)}, \\ \eta^3(\tau) \frac{\vartheta_1(\theta_1 + \theta_2|\tau)}{\vartheta_4(\theta_1|\tau)\vartheta_4(\theta_2|\tau)} &= -\frac{\sqrt{2}}{4} h e^{-2\eta_2 \omega_2 \theta_1 \theta_2} \frac{\sigma(t_1 + t_2)}{\sigma_3(t_1)\sigma_3(t_2)}.\end{aligned}\tag{5.17}$$

Using these expressions, we can obtain an explicit answer for the Donaldson-Witten function of product ruled surfaces in the chamber where the volume of F_g is small. For $2\lambda_0 = [\mathbb{C}P^1]$, we have to add the expression for the chamber where the volume of $[\mathbb{C}P^1]$ is small, and the infinite sum of wall-crossing terms. When the volume of $[\mathbb{C}P^1]$ is small, we have to use (5.4) but with the $-i \csc((S, [\mathbb{C}P^1])/2h)$ function as in (5.6). We then obtain

$$\begin{aligned}Z_{DW}^{[CP^1]} &= -2^{(7g+1)/2} (-i\pi)^g \left[h^{2g-1} f_2(q)^{-1} e^{(2p + \frac{S^2}{3})u} \right. \\ &\quad \cdot \left. \int_{\mathbf{T}^{b_1}} \exp \left\{ \frac{i\sqrt{2}}{24\pi} M(q)(S, [\mathbb{C}P^1])\Omega + Q(q)\gamma^\# \right\} \cdot \frac{\sigma_3(t_1 + t_2)}{\sigma(t_1)\sigma_3(t_2)} \right]_{q^0},\end{aligned}\tag{5.18}$$

where

$$M(q) = \frac{\vartheta_2^4 + \vartheta_3^4}{\vartheta_4^8}.\tag{5.19}$$

For $2\lambda_0 = [\mathbb{C}P^1] + [F_g]$ we obtain:

$$\begin{aligned}Z_{DW}^{[CP^1] + [F_g]} &= \frac{\sqrt{2}}{8} 2^{7g/2} (-i\pi)^g \left[h^{2g-1} f_2(q)^{-1} e^{(2p + \frac{S^2}{3})u} \right. \\ &\quad \cdot \left. \int_{\mathbf{T}^{b_1}} \exp \left\{ \frac{i\sqrt{2}}{24\pi} M(q)(S, [\mathbb{C}P^1])\Omega + Q(q)\gamma^\# \right\} \cdot \frac{\sigma(t_1 + t_2)}{\sigma_3(t_1)\sigma_3(t_2)} \right]_{q^0}.\end{aligned}\tag{5.20}$$

As a check of (5.20), one can easily derive, using the expansion of the σ functions, the result

$$\mathcal{D}^{[CP^1] + [F_1]}((I(\mathbb{C}P^1))^2) = -2,\tag{5.21}$$

where $\mathcal{D}^{[CP^1] + [F_1]}$ denotes the Donaldson invariant in the chamber $\text{vol}(F_1) \rightarrow 0$. This agrees with the computation in [2].

Recently the proof of the Atiyah conjecture on the relation of symplectic and instanton Floer homology has been completed [24][25]. It is possible that the above expressions can be used to give another proof of this conjecture. In fact,

$$Z_{DW}^{([CP^1],[CP^1]+[F_g])} = Z_{DW}^{[CP^1]} + Z_{DW}^{[CP^1]+[F_g]} \quad (5.22)$$

is a generating function for the Gromov-Witten invariants of the moduli space of flat connections [2][3][24][25]. In the simple case $g = 1$, one finds that

$$\mathcal{D}^{([CP^1],[CP^1]+[F_1])}(1) = -1, \quad \mathcal{D}^{([CP^1],[CP^1]+[F_1])}(\mathcal{O}) = 2, \quad (5.23)$$

where $\mathcal{O}$ is the zero-observable. Taking into account that the generator of the quantum cohomology of the moduli space of flat connections on F_1 (or of the Floer cohomology of $F_1 \times \mathbf{S}^1$) is given by $\beta = -4\mathcal{O}$, we obtain the relation $\beta = 8$, which is the first quantum correction to the classical cohomology ring in genus one.

Acknowledgements

We would like to thank Y. Ruan and E. Witten for some discussions. We would also like to thank V. Muñoz and Tianjun Li for very useful and clarifying correspondence. This work is supported by DOE grant DE-FG02-92ER40704.

References

- [1] *The Floer Memorial Volume*, H. Hofer et. al. eds., Birkäuser 1995.
- [2] S. Donaldson, “Floer homology and algebraic geometry,” in *Vector bundles in algebraic geometry*, N.J. Hitchin et. al. eds. Cambridge 1995.
- [3] M. Bershadsky, A. Johansen, V. Sadov, and C. Vafa, “Topological Reduction of 4D SYM to 2D σ -Models,” hep-th/9501096; Nucl. Phys. **B448** (1995) 166.
- [4] J. Morgan and T. Mrowka, “A note on Donaldson’s polynomial invariants,” Int. Math. Res. Not. **10** (1992) 223.
- [5] E. Witten, “Topological Quantum Field Theory,” Commun. Math. Phys. **117** (1988) 353.
- [6] E. Witten, “Supersymmetric Yang-Mills theory on a four-manifold,” hep-th/9403193; J. Math. Phys. **35** (1994) 5101.
- [7] E. Witten, “Monopoles and four-manifolds,” hep-th/9411102; Math. Res. Letters **1** (1994) 769.
- [8] E. Witten, “On S -duality in abelian gauge theory,” hep-th/9505186; Selecta Mathematica **1** (1995) 383.
- [9] G. Moore and E. Witten, “Integration over the u -plane in Donaldson theory,” hep-th/9709193.
- [10] A. Losev, N. Nekrasov, and S. Shatashvili, “Issues in topological gauge theory,” hep-th/9711108; “Testing Seiberg-Witten solution,” hep-th/9801061.
- [11] M. Mariño and G. Moore, “Integrating over the Coulomb branch in $\mathcal{N} = 2$ gauge theory,” hep-th/9712062.
- [12] M. Mariño and G. Moore, “The Donaldson-Witten function for gauge groups of rank larger than one,” hep-th/9802185.
- [13] Kanehisa Takasaki, “Integrable Hierarchies and Contact Terms in u -plane Integrals of Topologically Twisted Supersymmetric Gauge Theories,” hep-th/9803217
- [14] E. Verlinde, “Global aspects of electric-magnetic duality,” hep-th/9506011; Nucl. Phys. **B455** (1995) 211.
- [15] S.K. Donaldson and P.B. Kronheimer, *The Geometry of Four-Manifolds*, Clarendon Press, Oxford, 1990.
- [16] A. Gorsky, A. Marshakov, A. Mironov and A. Morozov, “RG equations from Whitham hierarchy,” hep-th/9802007.
- [17] V. Muñoz, “Wall-crossing formulae for algebraic surfaces with $q > 0$,” alg-geom/9709002. ■
- [18] L. Göttsche, “Modular forms and Donaldson invariants for 4-manifolds with $b_+ = 1$,” alg-geom/9506018; J. Am. Math. Soc. **9** (1996) 827.
- [19] L. Göttsche and D. Zagier, “Jacobi forms and the structure of Donaldson invariants for 4-manifolds with $b_+ = 1$,” alg-geom/9612020.
- [20] T.J. Li and A. Liu, “General wall-crossing formula,” Math. Res. Lett. **2** (1995) 797.

- [21] C. Okonek and A. Teleman, “Seiberg-Witten invariants for manifolds with $b_2^+ = 1$, and the universal wall-crossing formula,” alg-geom/9603003; Int. J. Math. **7** (1996) 811.
- [22] A. Weil, *Elliptic Functions according to Eisenstein and Kronecker*, Springer-Verlag, 1976
- [23] A. Polishchuk, “Massey and Fukaya products on elliptic curves,” alg-geom/9803017.
- [24] V. Muñoz, “Ring structure of the Floer cohomology of $\Sigma \times \mathbf{S}^1$,” dg-ga/9710029.
- [25] V. Muñoz, “Quantum cohomology of the moduli space of stable bundles over a Riemann surface,” alg-geom/9711030.
- [26] R. Fintushel and R.J. Stern, “The blowup formula for Donaldson invariants,” alg-geom/9405002; Ann. Math. **143** (1996) 529.